



Abstract: 5746

Scientific Abstracts &gt; Emerging Technology

## Exploring Mobile Health Solutions for Pain Management and Assessment of Spinal Joint Injection Efficacy

Ted Miclau, Matthew Smuck, Brandon Goenawan, Conor O'Neill, Lyndly Tamura, Peter Wu, Patricia Zheng, Cara Prideaux  
Mayo Clinic

### Introduction

Tracking outcomes in spine interventions is costly and time-consuming, hindering research progress. One recent randomized controlled trial examining lumbar epidural steroid injections required a budget exceeding two million dollars. Significant portions of research budgets are allocated to hiring personnel for patient recruitment and data collection. As 77% of Americans own smartphones, mobile health (mHealth) applications offer solutions to streamline the research process. These applications allow for remote enrollment and direct engagement with patients, reducing the need to hire research personnel to recruit patients and gather their follow-up outcomes. Studies demonstrate that utilization of mHealth applications may promote more effective post-operative care, as evidenced by an association with reduced emergency department visits and hospital readmissions. However, the existing literature on mHealth interventions for pain management is limited and heterogeneous.

Our study assessed the feasibility of a custom mHealth application designed to track longitudinal pain outcomes after various spine injections. Specifically, we tested the recruitment and retention capabilities of this custom mobile application paired with several recruitment strategies.

### Materials and Methods

IRB approval was obtained for this prospective observational study, which approached and enrolled a convenience sample of eligible patients scheduled for spine injections at an academic outpatient surgical center. After screening for inclusion and exclusion criteria, participants were recruited in the pre-injection environment through one of three modalities: in-person research assistant (RA), remote RA, and in-person clinician recruitment. Upon enrollment, a custom-built web-based mHealth application prompted patients to respond to post-injection surveys at pre-defined intervals based on injection type (e.g., epidural steroid injection (ESI), facet joint injection, medial branch block (MBB), and radiofrequency ablation (RFA)). The rates of successful study enrollment (i.e., percentage of patients enrolled from total number approached) and retention (i.e., percentage of requested pain surveys completed by patients) were compared between the different types of patient recruitment (i.e., clinician versus research assistant; remote versus in-person).

### Results/Case Report

Eighty-nine patients were approached for study enrollment. Enrollment success was 95.5%, 85.7%, and 25.6%

amongst the in-person clinician (N=22), in-person RA (N=28), and remote RA (N=39) recruitment groups, respectively. The percentage of completed surveys was 73.9%, 77.3%, and 74.3% amongst the in-person clinician (N=21), in-person RA (N=21), and remote RA recruitment groups (N=9), respectively. Enrollment differed significantly between groups (Fisher's exact test,  $p < .001$ ), with remote recruitment exhibiting less success than in-person recruitment. One-way ANOVA showed no significant differences in the percentage of surveys completed by patients between groups ( $p = .915$ ).

## Discussion

The use of an mHealth application to track post-spine injection pain outcomes appears feasible. Varied recruitment methods paired with the use of a mobile application to track post-spine injection pain outcomes yielded high enrollment and retention rates. In-person recruitment was more effective than remote recruitment, and clinician recruitment may be more effective than non-clinician recruitment. Recruitment approach was not associated with significant differences in post-enrollment survey completion rates. We continue to evaluate how different recruitment strategies can pair with a mobile application to alleviate research burdens and optimize patient enrollment and retention in the study of spine injection outcomes.

## References

1. Friedly JL, Comstock BA, Turner JA, et al. A Randomized Trial of Epidural Glucocorticoid Injections for Spinal Stenosis. *New England Journal of Medicine*. 2014;371(1). doi:10.1056/nejmoa1313265
2. Ghahreman A, Ferch R, Bogduk N. The efficacy of transforaminal injection of steroids for the treatment of lumbar radicular pain. *Pain Medicine (United States)*. 2010;11(8). doi:10.1111/j.1526-4637.2010.00908.x
3. Manchikanti L, Pampati V, Bakhit CE, et al. Manchikanti et al • Effectiveness of Lumbar Facet Joint Nerve Blocks Effectiveness of Lumbar Facet Joint Nerve Blocks in Chronic Low Back Pain: A Randomized Clinical Trial. Vol 4.; 2001.
4. Gliklich RE, Dreyer NA, Leavy MB. *Registries for Evaluating Patient Outcomes: A User's Guide*. Vol 1.; 2014.
5. Dawes J, Lin AY, Varghese C, Russell MM, Lin AY. Mobile health technology for remote home monitoring after surgery: a meta-analysis. *British Journal of Surgery*. 2021;108(11). doi:10.1093/bjs/zxab323
6. Piguet V, Mazouri S, Rochat J, Ehrler F, Cedraschi C, Luthy C. Mobile applications and self-management of chronic pain. *Rev Med Suisse*. 2020;16(700).
7. Koumpouros Y, Georgoulas A. Pain Management Mobile Applications: A Systematic Review of Commercial and Research Efforts. *Sensors*. 2023;23(15). doi:10.3390/s23156965
8. Almeida AF, Rocha NP, Silva AG. Methodological quality of manuscripts reporting on the usability of mobile applications for pain assessment and management: A systematic review. *Int J Environ Res Public Health*. 2020;17(3). doi:10.3390/ijerph17030785

## Disclosures

No

Tables / Images

□